

Lampiran Hasil Analisis Data

a. Karakteristik Subjek Penelitian

JK_balita

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Laki-laki	74	49.3	49.3	49.3
	Perempuan	76	50.7	50.7	100.0
	Total	150	100.0	100.0	

Umur_balita

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	12-23 bulan	46	30.7	30.7	30.7
	24-35 bulan	31	20.7	20.7	51.3
	36-47 bulan	40	26.7	26.7	78.0
	48-59 bulan	33	22.0	22.0	100.0
	Total	150	100.0	100.0	

JK_orbu

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Laki-laki	3	2.0	2.0	2.0
	Perempuan	147	98.0	98.0	100.0
	Total	150	100.0	100.0	

Umur_orbu1

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	15-24 tahun	13	8.7	8.7	8.7
	25-34 tahun	88	58.7	58.7	67.3
	35-44 tahun	48	32.0	32.0	99.3
	55-60 tahun	1	.7	.7	100.0
	Total	150	100.0	100.0	

Pekerjaan

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Buruh	6	4.0	4.0	4.0
	Guru	6	4.0	4.0	8.0
	Ibu Rumah Tangga	93	62.0	62.0	70.0
	Karyawan Swasta	31	20.7	20.7	90.7
	Pedagang	2	1.3	1.3	92.0
	Penjahit	4	2.7	2.7	94.7
	Perawat	2	1.3	1.3	96.0
	TU	1	.7	.7	96.7
	Wiraswasta	5	3.3	3.3	100.0
	Total	150	100.0	100.0	

Penghasilan ortu

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid < UMR	99	66.0	66.0	66.0
≥ UMR	51	34.0	34.0	100.0
Total	150	100.0	100.0	

Pend terakhir

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid Tamat SD	6	4.0	4.0	4.0
Tamat SMP	40	26.7	26.7	30.7
Tamat SMA/SMK	71	47.3	47.3	78.0
Diploma D1/D3	8	5.3	5.3	83.3
Sarjana (D4/S1)	25	16.7	16.7	100.0
Total	150	100.0	100.0	

b. Analisis Univariat

Status gizi

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Gizi Kurang (Wasting)	50	33.3	33.3	33.3
	Gizi Normal	100	66.7	66.7	100.0
	Total	150	100.0	100.0	

ASI Eksklusif

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Tidak ASI Eksklusif	77	51.3	51.3	51.3
	ASI Eksklusif	73	48.7	48.7	100.0
	Total	150	100.0	100.0	

Diare

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Ya	41	27.3	27.3	27.3
	Tidak	109	72.7	72.7	100.0
	Total	150	100.0	100.0	

ISPA

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Ya	103	68.7	68.7	68.7
	Tidak	47	31.3	31.3	100.0
	Total	150	100.0	100.0	

TBC

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Ya	3	2.0	2.0	2.0
	Tidak	147	98.0	98.0	100.0
	Total	150	100.0	100.0	

c. Analisis Bivariat

Crosstab

			Status_gizi		Total
			Gizi Kurang (Wasting)	Gizi Normal	
ASI_Eksklusif	Tidak ASI Eksklusif	Count	33	44	77
		Expected Count	25.7	51.3	77.0
		% within ASI_Eksklusif	42.9%	57.1%	100.0%
		% within Status_gizi	66.0%	44.0%	51.3%
		% of Total	22.0%	29.3%	51.3%
ASI Eksklusif	ASI Eksklusif	Count	17	56	73
		Expected Count	24.3	48.7	73.0
		% within ASI_Eksklusif	23.3%	76.7%	100.0%
		% within Status_gizi	34.0%	56.0%	48.7%
		% of Total	11.3%	37.3%	48.7%
Total		Count	50	100	150
		Expected Count	50.0	100.0	150.0
		% within ASI_Eksklusif	33.3%	66.7%	100.0%
		% within Status_gizi	100.0%	100.0%	100.0%
		% of Total	33.3%	66.7%	100.0%

Chi-Square Tests

	Value	df	Asymptotic Significance (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	6.458 ^a	1	.011	.015	.009
Continuity Correction ^b	5.607	1	.018		
Likelihood Ratio	6.548	1	.011		
Fisher's Exact Test					
Linear-by-Linear Association	6.415	1	.011		
N of Valid Cases	150				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 24,33.

b. Computed only for a 2x2 table

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for ASI_Eksklusif (Tidak ASI Eksklusif / ASI Eksklusif)	2.471	1.220	5.005
For cohort Status_gizi = Gizi Kurang (Wasting)	1.840	1.128	3.003
For cohort Status_gizi = Gizi Normal	.745	.591	.939
N of Valid Cases	150		

Crosstab

			Status_gizi		Total
			Gizi Kurang (Wasting)	Gizi Normal	
Diare	Ya	Count	17	24	41
		Expected Count	13.7	27.3	41.0
		% within Diare	41.5%	58.5%	100.0%
		% within Status_gizi	34.0%	24.0%	27.3%
		% of Total	11.3%	16.0%	27.3%
	Tidak	Count	33	76	109
		Expected Count	36.3	72.7	109.0
		% within Diare	30.3%	69.7%	100.0%
		% within Status_gizi	66.0%	76.0%	72.7%
		% of Total	22.0%	50.7%	72.7%
Total	Count		50	100	150
	Expected Count		50.0	100.0	150.0
	% within Diare		33.3%	66.7%	100.0%
	% within Status_gizi		100.0%	100.0%	100.0%
	% of Total		33.3%	66.7%	100.0%

Chi-Square Tests

	Value	df	Asymptotic Significance (2- sided)	Exact Sig. (2- sided)	Exact Sig. (1- sided)
Pearson Chi-Square	1.678 ^a	1	.195	.244	.136
Continuity Correction ^b	1.213	1	.271		
Likelihood Ratio	1.644	1	.200		
Fisher's Exact Test					
Linear-by-Linear Association	1.667	1	.197		
N of Valid Cases	150				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 13,67.

b. Computed only for a 2x2 table

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for Diare (Ya / Tidak)	1.631	.776	3.432
For cohort Status_gizi = Gizi Kurang (Wasting)	1.370	.863	2.174
For cohort Status_gizi = Gizi Normal	.840	.631	1.117
N of Valid Cases	150		

Crosstab

			Status_gizi		Total
			Gizi Kurang (Wasting)	Gizi Normal	
ISPA	Ya	Count	35	68	103
		Expected Count	34.3	68.7	103.0
		% within ISPA	34.0%	66.0%	100.0%
		% within Status_gizi	70.0%	68.0%	68.7%
		% of Total	23.3%	45.3%	68.7%
	Tidak	Count	15	32	47
		Expected Count	15.7	31.3	47.0
		% within ISPA	31.9%	68.1%	100.0%
		% within Status_gizi	30.0%	32.0%	31.3%
		% of Total	10.0%	21.3%	31.3%
Total	Count		50	100	150
	Expected Count		50.0	100.0	150.0
	% within ISPA		33.3%	66.7%	100.0%
	% within Status_gizi		100.0%	100.0%	100.0%
	% of Total		33.3%	66.7%	100.0%

Chi-Square Tests

	Value	df	Asymptotic Significance (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	.062 ^a	1	.803	.854	.478
Continuity Correction ^b	.004	1	.950		
Likelihood Ratio	.062	1	.803		
Fisher's Exact Test					
Linear-by-Linear Association	.062	1	.804		
N of Valid Cases	150				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 15,67.

b. Computed only for a 2x2 table

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for ISPA (Ya / Tidak)	1.098	.526	2.293
For cohort Status_gizi = Gizi Kurang (Wasting)	1.065	.648	1.750
For cohort Status_gizi = Gizi Normal	.970	.763	1.232
N of Valid Cases	150		

Crosstab

			Status_gizi		Total
			Gizi Kurang (Wasting)	Gizi Normal	
TBC	Ya	Count	1	2	3
		Expected Count	1.0	2.0	3.0
		% within TBC	33.3%	66.7%	100.0%
		% within Status_gizi	2.0%	2.0%	2.0%
		% of Total	0.7%	1.3%	2.0%
	Tidak	Count	49	98	147
		Expected Count	49.0	98.0	147.0
		% within TBC	33.3%	66.7%	100.0%
		% within Status_gizi	98.0%	98.0%	98.0%
		% of Total	32.7%	65.3%	98.0%
Total	Count		50	100	150
	Expected Count		50.0	100.0	150.0
	% within TBC		33.3%	66.7%	100.0%
	% within Status_gizi		100.0%	100.0%	100.0%
	% of Total		33.3%	66.7%	100.0%

Chi-Square Tests

	Value	df	Asymptotic Significance (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	.000 ^a	1	1.000	1.000	.742
Continuity Correction ^b	.000	1	1.000		
Likelihood Ratio	.000	1	1.000		
Fisher's Exact Test					
Linear-by-Linear Association	.000	1	1.000		
N of Valid Cases	150				

a. 2 cells (50,0%) have expected count less than 5. The minimum expected count is 1,00.

b. Computed only for a 2x2 table

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for TBC (Ya / Tidak)	1.000	.088	11.300
For cohort Status_gizi = Gizi Kurang (Wasting)	1.000	.199	5.036
For cohort Status_gizi = Gizi Normal	1.000	.446	2.244
N of Valid Cases	150		